


## Free PSA: Improving the discrimination between BPH and prostatic carcinoma

"The ratio of free/total PSA improves the discrimination between BPH and prostatic CA, particularly in the intermediate or "grey zone" range of 3-10 ng/ml of total PSA."



Total PSA (prostate specific antigen) assays have been widely used in recent years in the evaluation of patients with prostatic carcinoma (CA) and benign prostatic hypertrophy (BPH).

More recently the ratio of free PSA to complexed PSA has been found by several groups to be superior in differentiating raised levels of total PSA due to prostatic CA from raised levels due to BPH.

PSA, which is a serine protease with chymotrypsinogen-like activity, occurs in different molecular forms in serum. Most of the immunodetectable PSA found in serum is complexed to a serine protease inhibitor called alpha-1- antichymotrypsin (PSA-ACT). Free PSA or uncomplexed PSA is the other immunodetectable form in serum.

Several groups have now found that the proportion of free PSA in serum is significantly higher in patients with BPH than in patients with prostate cancer.

**The ratio of free/total PSA improves the discrimination between BPH and prostatic CA, particularly in the intermediate or "grey zone" range of 3-10 ng/ml of total PSA.**


In one of the largest series [Apoorva et al (Oesterling) Urology, 49: 19, 1997] a total PSA level of between 3-4 ng/ml with a free to total ratio of less than 0.19 is regarded as an indication for ultrasound biopsy. In this situation the sensitivity is 90% and the specificity 48%, indicating that 48% of unnecessary biopsies would be eliminated and 10% of cancers would be missed. Lowering the cut point to 0.15 decreases sensitivity to 80% and increases specificity to 62%.

At levels of 4.1-10 ng/ml using a cut off ratio of 0.24 or less, the sensitivity is 95%, and the specificity is 13%, i.e. 13% of unnecessary biopsies would be eliminated and 5% of cancers missed. Using a cut off point of 0.22, sensitivity was 90% and specificity 19%.

In two other series (Partin et al, Urology 48:55,1996 and Alivizatos et al, Urology, 48:71,1996) using a cut off point of 0.20 in the 4-10 ng/ml total PSA range, sensitivities were 95% and 77% respectively and specificities 29% and 65% respectively. From this it can be seen that the results vary in different series.

The free/total cut off point is also dependent on the clinician, depending on whether he desires increased sensitivity of detection of cancer (i.e. raising the cut point), or increased specificity (i.e. decreasing the biopsy rate) by lowering the cut off point.

These recommendations are summarized in a testing algorithm for PSA recommended by Oesterling. July 2003



**PATHOLOGY IS OUR CORE BUSINESS. WELCOME TO THE LANCET EXPERIENCE**

*Material from Lancet SA Newsletter - Compiled by Dr. Juanita Smit - January 2011*

### Branches/Service points Countrywide

**LANCET- MOMBASA**

Ground Floor within Ocean Medical Centre opposite Law Courts .  
Tel: 0721 143 766/ 0736 489282

**LANCET- KISUMU**

2nd floor WEDCO Centre on Oginga Odinga Street  
Tel: 0726 838773/ 0735 457665

**LANCET- THIKA**

3rd floor Thika Arcade  
Tel: 020262 2633/ 0717414684

**LANCET - NELSON AWORI**

Professor Nelson Awori Centre, 2nd floor next to The Nairobi Hospital  
Tel: 0726 839341

**LANCET- EASTLEIGH**

Alliance Medical Centre Madina Shopping Mall 2nd floor opp. Roundabout of Gen. Waruinge Rd/ 1st Avenue Tel: 0717 414682

**LANCET- GA HOUSE**

3rd floor General Accident House within Zenith Medical Centre, Ralphe Banche Road, Upper Hill - Nairobi  
Tel: 0726 995 860

**LANCET- SEARCH LAB**

Hurlingham Medical Plaza, Ground floor adjacent to the Nairobi Women's Hospital on Argwings Kodhek Road

**LANCET- BURUBURU**

Buruburu service point behind Misora Tel: 0717414708

**LANCET- PARKLANDS**

Parklands MediPlaza 2nd floor within The International Clinic (opposite the Aga Khan Hospital) Tel: 0729 909409

**LANCET- CBD**

Opening Soon